

## FORMS OF IMMUNE AUGMENTATION

1. FREE ANTIBODY + PEPTIDE





# 2. HETEROCONJUGATE (ANTIBODY-PEPTIDE)



# 3. BIFUNCTIONAL mAb/LIGAND CONJUGATE




FIGURE 2:







FIGURE 3:








ALLO PRE-OK


₩ ALLO POST-(

(CG) HOURS PO


DOSE



ANTI-KLH ANTIBODIES IN MICE TREATED
WITH KLH USING PBS, CFA, OR ANTI-CD3 AS IMMUNOADJUVANTS



# **ALLORESPONSE TO SEB**



in C3H mice at 18 hr.








FIGURE 8:


# PROLIFERATIVE RESPONSE TO SEB



in C3H mice at 18 hr. using lymph node


# EXPANSION OF % 8+ CELLS IN SEB-TREATED MICE






STASIK COUNTS

## Absolute Counts





NL HAM Ig

ANTI-CD3 (250ug)

ANTI-CD3 (25ug)

20

0

C

ರ

5


20

25


30


DAYS POST TREATMENT

## ANTI-1005/45 TITER IN C57BL/10 MICE GIVEN 1005/45-KLH SC OR IP AND 145-2C11




ANTI-CD3 VS. F(ab')2 ANTI-CD3 x ANTI-CD4 HETEROCONJUGATE IN VIVO TREATMENT WITH mAbs:






### FIGURE 17:


### TWO MODELS OF TB COLLABORATION





MHC RESTRICTED AG RECOGNITION



mAb-A(ANTI-CD3) PEPTIDE










FIG. 3




FIG. 4A






FIG. 4B









FIG. 6






FIG. 8A




FIG. 8B




FIG. 8C





FIG. 8D






FIG. 11




FIG. 12




FIG. 13




FIG. 14





FIG. 15








FIG. 16B